

ON ENS PORTA EL DSM-5? LA DEPRESSIÓ AL DSM-5, QUÈ SOLUCIONA?

VÍCTOR PÉREZ SOLA

Servei de Psiquiatria

Hospital de Sant Pau, Barcelona

Institut d'Investigació Biomèdica Sant Pau (IIB-Sant Pau)

Centro de Investigación Biomédica en Red de Salud

Mental (CIBERSAM)



cibersam
Centro de Investigación Biomédica en Red de Salud Mental


ib **santpau**
Hospital de Sant Pau

HOSPITAL DE LA
SANTA CREU I
SANT PAU
CIBERSAM

Essay

A Comparison of DSM-IV and DSM-5 Panel Members' Financial Associations with Industry: A Pernicious Problem Persists

Lisa Cosgrove^{1,2*}, Sheldon Krinsky³



Contents

DSM-5 Classification	xiii
Preface	xii

Section I DSM-5 Basics

Introduction	5
Use of the Manual	19
Cautionary Statement for Forensic Use of DSM-5	25

Section II Diagnostic Criteria and Codes

Neurodevelopmental Disorders	31
Schizophrenia Spectrum and Other Psychotic Disorders	87
Bipolar and Related Disorders	123
Depressive Disorders	155
Anxiety Disorders	189
Obsessive-Compulsive and Related Disorders	235
Trauma- and Stressor-Related Disorders	265
Dissociative Disorders	291
Somatic Symptom and Related Disorders	309
Feeding and Eating Disorders	329
Elimination Disorders	355
Sleep-Wake Disorders	361
Sexual Dysfunctions	423
Gender Dysphoria	451



Contents

DSM-5 Classification	xiii
Preface	xii

Section I DSM-5 Basics

Introduction	5
Use of the Manual	19
Cautionary Statement for Forensic Use of DSM-5	25

Section II Diagnostic Criteria and Codes

Neurodevelopmental Disorders	31
Schizophrenia Spectrum and Other Psychotic Disorders	87
Bipolar and Related Disorders	123
Depressive Disorders	155
Anxiety Disorders	189
Obsessive-Compulsive and Related Disorders	235
Trauma- and Stressor-Related Disorders	265
Dissociative Disorders	291
Somatic Symptom and Related Disorders	309
Feeding and Eating Disorders	329
Elimination Disorders	355
Sleep-Wake Disorders	361
Sexual Dysfunctions	423
Gender Dysphoria	451

DIAGNOSTIC AND STATISTICAL
MANUAL OF
MENTAL DISORDERS

FIFTH EDITION

DSM-5

AMERICAN PSYCHIATRIC ASSOCIATION

Trastorno Bipolar



Con la intención de mejorar la precisión del diagnóstico y facilitar la detección temprana en el ámbito clínico, el Criterio

A para episodios maníacos y hipomaníacos ahora incluye un énfasis en la evolución de la actividad y la energía, así como el estado de ánimo.

Trastorno Bipolar



Se ha eliminado el diagnóstico DSM-IV de trastorno bipolar I, episodio mixto que requería que la persona cumpliera simultáneamente todos los criterios para la manía y depresión mayor.

Se propone un nuevo especificador:

Con “**características mixtas**” que se puede aplicar a los episodios de manía o hipomanía cuando hay síntomas depresivos y en episodios depresivos en el contexto de un trastorno depresivo mayor o trastorno bipolar cuando las características de la manía / hipomanía están presentes.

Trastorno Bipolar



With mixed features: The mixed features specifier can apply to the current manic, hypomanic, or depressive episode in bipolar I or bipolar II disorder.

Manic or hypomanic episode, with mixed features:

- A. Full criteria are met for a manic episode or hypomanic episode, and at least three of the following symptoms are present during the majority of days of the current or most recent episode of mania or hypomania:
1. Prominent dysphoria or depressed mood as indicated by either subjective report (e.g., feels sad or empty) or observation made by others (e.g., appears tearful).
 2. Diminished interest or pleasure in all, or almost all, activities (as indicated by either subjective account or observation made by others).
 3. Psychomotor retardation nearly every day (observable by others; not merely subjective feelings of being slowed down).
 4. Fatigue or loss of energy.
 5. Feelings of worthlessness or excessive or inappropriate guilt (not merely self-reproach or guilt about being sick).
 6. Recurrent thoughts of death (not just fear of dying), recurrent suicidal ideation without a specific plan, or a suicide attempt or a specific plan for committing suicide.
- B. Mixed symptoms are observable by others and represent a change from the person's usual behavior.
- C. For individuals whose symptoms meet full episode criteria for both mania and depression simultaneously, the diagnosis should be manic episode, with mixed features, due to the marked impairment and clinical severity of full mania.
- D. The mixed symptoms are not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication, other treatment).

Trastorno Bipolar



Depressive episode, with mixed features:

- A. Full criteria are met for a major depressive episode, and at least three of the following manic/hypomanic symptoms are present during the majority of days of the current or most recent episode of depression:
1. Elevated, expansive mood.
 2. Inflated self-esteem or grandiosity.
 3. More talkative than usual or pressure to keep talking.
 4. Flight of ideas or subjective experience that thoughts are racing.
 5. Increase in energy or goal-directed activity (either socially, at work or school, or sexually).
 6. Increased or excessive involvement in activities that have a high potential for painful consequences (e.g., engaging in unrestrained buying sprees, sexual indiscretions, or foolish business investments).
 7. Decreased need for sleep (feeling rested despite sleeping less than usual; to be contrasted with insomnia).
- B. Mixed symptoms are observable by others and represent a change from the person's usual behavior.
- C. For individuals whose symptoms meet full episode criteria for both mania and depression simultaneously, the diagnosis should be manic episode, with mixed features.
- D. The mixed symptoms are not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication, or other treatment).

Note: Mixed features associated with a major depressive episode have been found to be a significant risk factor for the development of bipolar I or bipolar II disorder. As a result, it is clinically useful to note the presence of this specifier for treatment planning and monitoring of response to treatment.

Trastorno Bipolar

Con “Sintomatología ansiosa”

Destinada a identificar a los pacientes con síntomas de ansiedad que no forman parte de los criterios diagnósticos



Specifiers for Bipolar and Related Disorders

Specify if:

With anxious distress: The presence of at least two of the following symptoms during the majority of days of the current or most recent episode of mania, hypomania, or depression:

1. Feeling keyed up or tense.
2. Feeling unusually restless.
3. Difficulty concentrating because of worry.
4. Fear that something awful may happen.
5. Feeling that the individual might lose control of himself or herself.

Specify current severity:

Mild: Two symptoms.

Moderate: Three symptoms.

Moderate-severe: Four or five symptoms.

Severe: Four or five symptoms with motor agitation.

Note: Anxious distress has been noted as a prominent feature of both bipolar and major depressive disorder in both primary care and specialty mental health settings. High levels of anxiety have been associated with higher suicide risk, longer duration of illness, and greater likelihood of treatment nonresponse. As a result, it is clinically useful to specify accurately the presence and severity levels of anxious distress for treatment planning and monitoring of response to treatment.

Trastorno Bipolar



Depressive Episodes With Short-Duration Hypomania

Proposed Criteria

Lifetime experience of at least one major depressive episode meeting the following criteria:

A. At least two lifetime episodes of hypomanic periods that involve the required criterion symptoms below but are of insufficient duration (at least 2 days but less than 4 consecutive days) to meet criteria for a hypomanic episode. The criterion symptoms are as follows:

- A. A distinct period of abnormally and persistently elevated, expansive, or irritable mood and abnormally and persistently increased goal-directed activity or energy.
- B. During the period of mood disturbance and increased energy and activity, three (or more) of the following symptoms have persisted (four if the mood is only irritable), represent a noticeable change from usual behavior, and have been present to a significant degree:
 1. Inflated self-esteem or grandiosity.
 2. Decreased need for sleep (e.g., feels rested after only 3 hours of sleep).
 3. More talkative than usual or pressured to keep talking.
 4. Flight of ideas or subjective experience that thoughts are racing.
 5. Distractibility (i.e., attention too easily drawn to unimportant or irrelevant external stimuli), as reported or observed.
 6. Increase in goal-directed activity (either socially, at work or school, or sexually) or psychomotor agitation.
 7. Excessive involvement in activities that have a high potential for painful consequences (e.g., the individual engages in unrestrained buying sprees, sexual indiscretions, or foolish business investments).

C. The episode is associated with an unequivocal change in functioning that is uncharacteristic of the individual when not symptomatic.

D. The disturbance in mood and the change in functioning are observable by others.

E. The episode is not severe enough to cause marked impairment in social or occupational functioning or to necessitate hospitalization. If there are psychotic features, the episode is, by definition, manic.

F. The episode is not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication or other treatment).

Trastorno Depresivo



DSM-5 contiene varios trastornos depresivos nuevos:

Trastorno de disregulación del estado de ánimo perturbador

Trastorno disfórico premenstrual.

Trastorno Depresivo



Trastorno de disregulación del estado de ánimo perturbador

Se propone con el objetivo de disminuir el potencial de sobrediagnóstico y sobretratamiento del trastorno bipolar en los niños.

Se incluyen los niños hasta la edad de 18 años que presentan irritabilidad persistente y frecuentes episodios de descontrol conductual extremo.

Trastorno de disregulación del estado de ánimo perturbador



Disruptive Mood Dysregulation Disorder

296.99 (F34.8)

Diagnostic Criteria

- A. Severe recurrent temper outbursts manifested verbally (e.g., verbal rages) and/or behaviorally (e.g., physical aggression toward people or property) that are grossly out of proportion in intensity or duration to the situation or provocation.
- B. The temper outbursts are inconsistent with developmental level.
- C. The temper outbursts occur, on average, three or more times per week.
- D. The mood between temper outbursts is persistently irritable or angry most of the day, nearly every day, and is observable by others (e.g., parents, teachers, peers).
- E. Criteria A–D have been present for 12 or more months. Throughout that time, the individual has not had a period lasting 3 or more consecutive months without all of the symptoms in Criteria A–D.
- F. Criteria A and D are present in at least two of three settings (i.e., at home, at school, with peers) and are severe in at least one of these.
- G. The diagnosis should not be made for the first time before age 6 years or after age 18 years.
- H. By history or observation, the age at onset of Criteria A–E is before 10 years.
- I. There has never been a distinct period lasting more than 1 day during which the full symptom criteria, except duration, for a manic or hypomanic episode have been met.
Note: Developmentally appropriate mood elevation, such as occurs in the context of a highly positive event or its anticipation, should not be considered as a symptom of mania or hypomania.

J. The behaviors do not occur exclusively during an episode of major depressive disorder and are not better explained by another mental disorder (e.g., autism spectrum disorder, posttraumatic stress disorder, separation anxiety disorder, persistent depressive disorder [dysthymia]).
Note: This diagnosis cannot coexist with oppositional defiant disorder, intermittent explosive disorder, or bipolar disorder, though it can coexist with others, including major depressive disorder, attention-deficit/hyperactivity disorder, conduct disorder, and substance use disorders. Individuals whose symptoms meet criteria for both disruptive mood dysregulation disorder and oppositional defiant disorder should only be given the diagnosis of disruptive mood dysregulation disorder. If an individual has ever experienced a manic or hypomanic episode, the diagnosis of disruptive mood dysregulation disorder should not be assigned.

K. The symptoms are not attributable to the physiological effects of a substance or to another medical or neurological condition.

Trastorno Depresivo



Trastorno disfórico premenstrual

Basado en pruebas científicas sólidas, el trastorno disfórico premenstrual se ha movido de DSM-IV Apéndice B, "Criterios y ejes propuestos para estudios posteriores" a un criterio principal en el DSM -5.

Trastorno disfórico premenstrual



Premenstrual Dysphoric Disorder

625.4 (N94.3)

Diagnostic Criteria

A. In the majority of menstrual cycles, at least five symptoms must be present in the final week before the onset of menses, start to improve within a few days after the onset of menses, and become minimal or absent in the week postmenses.

B. One (or more) of the following symptoms must be present:

1. Marked affective lability (e.g., mood swings, feeling suddenly sad or tearful, or increased sensitivity to rejection).
 2. Marked irritability or anger or increased interpersonal conflicts.
 3. Marked depressed mood, feelings of hopelessness, or self-deprecating thoughts.
 4. Marked anxiety, tension, and/or feelings of being keyed up or on edge.
- C.** One (or more) of the following symptoms must additionally be present, to reach a total of five symptoms when combined with symptoms from Criterion B above.
1. Decreased interest in usual activities (e.g., work, school, friends, hobbies).
 2. Subjective difficulty in concentration.
 3. Lethargy, easy fatigability, or marked lack of energy.
 4. Marked change in appetite; overeating; or specific food cravings.
 5. Hyperemia or insomnia.
 6. A sense of being overwhelmed or out of control.
 7. Physical symptoms, such as breast tenderness or swelling, joint or muscle pain, a sensation of "bloating," or weight gain.

Note: The symptoms in Criteria A–C must have been met for most menstrual cycles that occurred in the preceding year.

D. The symptoms are associated with clinically significant distress or interference with work, school, usual social activities, or relationships with others (e.g., avoidance of social activities; decreased productivity and efficiency at work, school, or home).

E. The disturbance is not merely an exacerbation of the symptoms of another disorder, such as major depressive disorder, panic disorder, persistent depressive disorder (dysthymia), or a personality disorder (although it may co-occur with any of these disorders).

F. Criterion A should be confirmed by prospective daily ratings during at least two symptomatic cycles. (**Note:** The diagnosis may be made provisionally prior to this confirmation.)

G. The symptoms are not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication, other treatment) or another medical condition (e.g., hypothyroidism).



Formas crónicas de depresión



La distimia del DSM-IV ahora cae bajo la categoría de trastorno depresivo persistente, que incluye el trastorno depresivo mayor crónico y el trastorno distímico anterior.

La incapacidad de encontrar diferencias científicamente significativas entre estas dos condiciones llevaron a su combinación con los especificadores incluye para identificar diferentes vías para el diagnóstico y para dar continuidad con el DSM-IV.

Formas crónicas de depresión



Persistent Depressive Disorder (Dysthymia)

300.4 (F34.1)

Diagnostic Criteria

This disorder represents a consolidation of DSM-IV-defined chronic major depressive disorder and dysthymic disorder.

A. Depressed mood for most of the day, for more days than not, as indicated by either subjective account or observation by others, for at least 2 years.

Note: In children and adolescents, mood can be irritable and duration must be at least 1 year.

B. Presence, while depressed, of two (or more) of the following:

1. Poor appetite or overeating.
2. Insomnia or hypersomnia.
3. Low energy or fatigue.
4. Low self-esteem.
5. Poor concentration or difficulty making decisions.
6. Feelings of hopelessness.

C. During the 2-year period (1 year for children or adolescents) of the disturbance, the individual has never been without the symptoms in Criteria A and B for more than 2 months at a time.

D. Criteria for a major depressive disorder may be continuously present for 2 years.

E. There has never been a manic episode or a hypomanic episode, and criteria have never been met for cyclothymic disorder.

F. The disturbance is not better explained by a persistent schizoaffective disorder, schizophrenia, delusional disorder, or other specified or unspecified schizophrenia spectrum and other psychotic disorder.

G. The symptoms are not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication) or another medical condition (e.g., hypothyroidism).

H. The symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.

Note: Because the criteria for a major depressive episode include four symptoms that are absent from the symptom list for persistent depressive disorder (dysthymia), a very limited

Formas crónicas de depresión



number of individuals will have depressive symptoms that have persisted longer than 2 years but will not meet criteria for persistent depressive disorder. If full criteria for a major depressive episode have been met at some point during the current episode of illness, they should be given a diagnosis of major depressive disorder. Otherwise, a diagnosis of other specified depressive disorder or unspecified depressive disorder is warranted.

Specify if:

- With anxious distress (p. 184)
- With mixed features (pp. 184–185)
- With melancholic features (p. 185)
- With atypical features (pp. 185–186)
- With mood-congruent psychotic features (p. 186)
- With mood-incongruent psychotic features (p. 186)
- With peripartum onset (pp. 186–187)

Specify if:

- In partial remission (p. 188)
- In full remission (p. 188)

Specify if:

- Early onset: If onset is before age 21 years.
- Late onset: If onset is at age 21 years or older.

Specify if (for most recent 2 years of persistent depressive disorder):

- With pure dysthymic syndrome: Full criteria for a major depressive episode have not been met in at least the preceding 2 years.
- With persistent major depressive episode: Full criteria for a major depressive episode have been met throughout the preceding 2-year period.
- With intermittent major depressive episodes, with current episode: Full criteria for a major depressive episode are currently met, but there have been periods of at least 8 weeks in at least the preceding 2 years with symptoms below the threshold for a full major depressive episode.
- With intermittent major depressive episodes, without current episode: Full criteria for a major depressive episode are not currently met, but there has been one or more major depressive episodes in at least the preceding 2 years.

Specify current severity:

- Mild (p. 188)
- Moderate (p. 188)
- Severe (p. 188)

Trastorno Depresivo Mayor



Si los síntomas centrales ni la duración requerida de al menos 2 semanas ha cambiado respecto al DSM-IV.

Criterio A para un episodio depresivo mayor en el DSM-5 es idéntica a la del DSM-IV.

Trastorno Depresivo Mayor



3. Significant weight loss when not dieting or weight gain (e.g., a change of more than 5% of body weight in a month), or decrease or increase in appetite nearly every day.
(**Note:** In children, consider failure to make expected weight gain.)
 4. Insomnia or hypersomnia nearly every day.
 5. Psychomotor agitation or retardation nearly every day (observable by others, not merely subjective feelings of restlessness or being slowed down).
 6. Fatigue or loss of energy nearly every day.
 7. Feelings of worthlessness or excessive or inappropriate guilt (which may be delusional) nearly every day (not merely self-reproach or guilt about being sick).
 8. Diminished ability to think or concentrate, or indecisiveness, nearly every day (either by subjective account or as observed by others).
 9. Recurrent thoughts of death (not just fear of dying), recurrent suicidal ideation without a specific plan, or a suicide attempt or a specific plan for committing suicide.
- B. The symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.
- C. The episode is not attributable to the physiological effects of a substance or to another medical condition.

Note: Criteria A–C represent a major depressive episode.

Note: Responses to a significant loss (e.g., bereavement, financial ruin, losses from a natural disaster, a serious medical illness or disability) may include the feelings of intense sadness, rumination about the loss, insomnia, poor appetite, and weight loss noted in Criterion A, which may resemble a depressive episode. Although such symptoms may be understandable or considered appropriate to the loss, the presence of a major depressive episode in addition to the normal response to a significant loss should also be carefully considered. This decision inevitably requires the exercise of clinical judgment based on the individual's history and the cultural norms for the expression of distress in the context of loss.¹

Trastorno Depresivo Mayor



¹In distinguishing grief from a major depressive episode (MDE), it is useful to consider that in grief the predominant affect is feelings of emptiness and loss, while in MDE it is persistent depressed mood and the inability to anticipate happiness or pleasure. The dysphoria in grief is likely to decrease in intensity over days to weeks and occurs in waves, the so-called pangs of grief. These waves tend to be associated with thoughts or reminders of the deceased. The depressed mood of MDE is more persistent and not tied to specific thoughts or preoccupations. The pain of grief may be accompanied by positive emotions and humor that are uncharacteristic of the pervasive unhappiness and misery characteristic of MDE. The thought content associated with grief generally features a preoccupation with thoughts and memories of the deceased, rather than the self-critical or pessimistic ruminations seen in MDE. In grief, self-esteem is generally preserved, whereas in MDE feelings of worthlessness and self-loathing are common. If self-derogatory ideation is present in grief, it typically involves perceived failings vis-à-vis the deceased (e.g., not visiting frequently enough, not telling the deceased how much he or she was loved). If a bereaved individual thinks about death and dying, such thoughts are generally focused on the deceased and possibly about "joining" the deceased, whereas in MDE such thoughts are focused on ending one's own life because of feeling worthless, undeserving of life, or unable to cope with the pain of depression.

Trastorno Depresivo Mayor



In recording the name of a diagnosis, terms should be listed in the following order: major depressive disorder, single or recurrent episode, severity/psychotic/remission specifiers, followed by as many of the following specifiers without codes that apply to the current episode.

Specify:

- With anxious distress** (p. 184)
- With mixed features** (pp. 184–185)
- With melancholic features** (p. 185)
- With atypical features** (pp. 185–186)
- With mood-congruent psychotic features** (p. 186)
- With mood-incongruent psychotic features** (p. 186)
- With catatonia** (p. 186). **Coding note:** Use additional code 293.89 (F06.1).
- With peripartum onset** (pp. 186–187)
- With seasonal pattern** (recurrent episode only) (pp. 187–188)

Trastorno Depresivo Mayor



Specifiers for Depressive Disorders

Specify if:

With anxious distress: Anxious distress is defined as the presence of at least two of the following symptoms during the majority of days of a major depressive episode or persistent depressive disorder (dysthymia):

1. Feeling keyed up or tense.
2. Feeling unusually restless.
3. Difficulty concentrating because of worry.
4. Fear that something awful may happen.
5. Feeling that the individual might lose control of himself or herself.

Specify current severity:

Mild: Two symptoms.

Moderate: Three symptoms.

Moderate-severe: Four or five symptoms.

Severe: Four or five symptoms and with motor agitation.

Note: Anxious distress has been noted as a prominent feature of both bipolar and major depressive disorder in both primary care and specialty mental health settings. High levels of anxiety have been associated with higher suicide risk, longer duration of illness, and greater likelihood of treatment nonresponse. As a result, it is clinically useful to specify accurately the presence and severity levels of anxious distress for treatment planning and monitoring of response to treatment.

Trastorno Depresivo Mayor



With mixed features:

- A. At least three of the following manic/hypomanic symptoms are present nearly every day during the majority of days of a major depressive episode:
1. Elevated, expansive mood.
 2. Inflated self-esteem or grandiosity.
 3. More talkative than usual or pressure to keep talking.
 4. Flight of ideas or subjective experience that thoughts are racing.
 5. Increase in energy or goal-directed activity (either socially, at work or school, or sexually).
 6. Increased or excessive involvement in activities that have a high potential for painful consequences (e.g., engaging in unrestrained buying sprees, sexual indiscretions, foolish business investments).
 7. Decreased need for sleep (feeling rested despite sleeping less than usual; to be contrasted with insomnia).
- B. Mixed symptoms are observable by others and represent a change from the person's usual behavior.
- C. For individuals whose symptoms meet full criteria for either mania or hypomania, the diagnosis should be bipolar I or bipolar II disorder.
- D. The mixed symptoms are not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication or other treatment).
- Note:** Mixed features associated with a major depressive episode have been found to be a significant risk factor for the development of bipolar I or bipolar II disorder. As a result, it is clinically useful to note the presence of this specifier for treatment planning and monitoring of response to treatment.

Trastornos Depresivos



Persistent Complex Bereavement Disorder

Proposed Criteria

- A. The individual experienced the death of someone with whom he or she had a close relationship.
- B. Since the death, at least one of the following symptoms is experienced on more days than not and to a clinically significant degree and has persisted for at least 12 months after the death in the case of bereaved adults and 6 months for bereaved children:
1. Persistent yearning/longing for the deceased. In young children, yearning may be expressed in play and behavior, including behaviors that reflect being separated from, and also reuniting with, a caregiver or other attachment figure.
 2. Intense sorrow and emotional pain in response to the death.
 3. Preoccupation with the deceased.
 4. Preoccupation with the circumstances of the death. In children, this preoccupation with the deceased may be expressed through the themes of play and behavior and may extend to preoccupation with possible death of others close to them.
- C. Since the death, at least six of the following symptoms are experienced on more days than not and to a clinically significant degree, and have persisted for at least 12 months after the death in the case of bereaved adults and 6 months for bereaved children:

Trastornos Depresivos



Reactive distress to the death

1. Marked difficulty accepting the death. In children, this is dependent on the child's capacity to comprehend the meaning and permanence of death.
2. Experiencing disbelief or emotional numbness over the loss.
3. Difficulty with positive reminiscing about the deceased.
4. Bitterness or anger related to the loss.
5. Maladaptive appraisals about oneself in relation to the deceased or the death (e.g., self-blame).
6. Excessive avoidance of reminders of the loss (e.g., avoidance of individuals, places, or situations associated with the deceased; in children, this may include avoidance of thoughts and feelings regarding the deceased).

Social/identity disruption

7. A desire to die in order to be with the deceased.
8. Difficulty trusting other individuals since the death.
9. Feeling alone or detached from other individuals since the death.
10. Feeling that life is meaningless or empty without the deceased, or the belief that one cannot function without the deceased.
11. Confusion about one's role in life, or a diminished sense of one's identity (e.g., feeling that a part of oneself died with the deceased).
12. Difficulty or reluctance to pursue interests since the loss or to plan for the future (e.g., friendships, activities).

D. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.

E. The bereavement reaction is out of proportion to or inconsistent with cultural, religious, or age-appropriate norms.

Specify if:

With traumatic bereavement: Bereavement due to homicide or suicide with persistent distressing preoccupations regarding the traumatic nature of the death (often in response to loss reminders), including the deceased's last moments, degree of suffering and mutilating injury, or the malicious or intentional nature of the death.

Trastornos Depresivos



Suicidal Behavior Disorder

Proposed Criteria

- A. Within the last 24 months, the individual has made a suicide attempt.
Note: A suicide attempt is a self-initiated sequence of behaviors by an individual who, at the time of initiation, expected that the set of actions would lead to his or her own death. The "time of initiation" is the time when a behavior took place that involved applying the method.)
- B. The act does not meet criteria for nonsuicidal self-injury—that is, it does not involve self-injury directed to the surface of the body undertaken to induce relief from a negative feeling/cognitive state or to achieve a positive mood state.
- C. The diagnosis is not applied to suicidal ideation or to preparatory acts.
- D. The act was not initiated during a state of delirium or confusion.
- E. The act was not undertaken solely for a political or religious objective.

Specify if:

Current: Not more than 12 months since the last attempt.

In early remission: 12–24 months since the last attempt.

Trastornos Depresivos



Nonsuicidal Self-Injury

Proposed Criteria

A. In the last year, the individual has, on 5 or more days, engaged in intentional self-inflicted damage to the surface of his or her body of a sort likely to induce bleeding, bruising, or pain (e.g., cutting, burning, stabbing, hitting, excessive rubbing), with the expectation that the injury will lead to only minor or moderate physical harm (i.e., there is no suicidal intent).

Note: The absence of suicidal intent has either been stated by the individual or can be inferred by the individual's repeated engagement in a behavior that the individual knows, or has learned, is not likely to result in death.

B. The individual engages in the self-injurious behavior with one or more of the following expectations:

1. To obtain relief from a negative feeling or cognitive state.
2. To resolve an interpersonal difficulty.
3. To induce a positive feeling state.

Note: The desired relief or response is experienced during or shortly after the self-injury, and the individual may display patterns of behavior suggesting a dependence on repeatedly engaging in it.

C. The intentional self-injury is associated with at least one of the following:

1. Interpersonal difficulties or negative feelings or thoughts, such as depression, anxiety, tension, anger, generalized distress, or self-criticism, occurring in the period immediately prior to the self-injurious act.
2. Prior to engaging in the act, a period of preoccupation with the intended behavior that is difficult to control.
3. Thinking about self-injury that occurs frequently, even when it is not acted upon.

D. The behavior is not socially sanctioned (e.g., body piercing, tattooing, part of a religious or cultural ritual) and is not restricted to picking a scab or nail biting.

E. The behavior or its consequences cause clinically significant distress or interference in interpersonal, academic, or other important areas of functioning.

F. The behavior does not occur exclusively during psychotic episodes, delirium, substance intoxication, or substance withdrawal. In individuals with a neurodevelopmental disorder, the behavior is not part of a pattern of repetitive stereotypies. The behavior is not better explained by another mental disorder or medical condition (e.g., psychotic disorder, autism spectrum disorder, intellectual disability, Lesch-Nyhan syndrome, stereotypic movement disorder with self-injury, trichotillomania [hair-pulling disorder], excoriation [skin-picking] disorder).